

PancrePlus™

Tablets and Powder

PancrePlus is a pancreatic enzyme concentrate of porcine origin. It is used as a digestive aid in replacement therapy where digestion of protein, carbohydrate, and fat is inadequate due to exocrine pancreatic insufficiency. Available in tablets or powder.

Product	Item#	Size
PancrePlus Tablets	501048	100 ct
	502048	500 ct
PancrePlus Powder	503048	4 oz
	504048	8 oz
	505048	12 oz







vetone[®] PancrePlus[™]

Tablets Pancreatic Enzyme Concentrate

For use in animals only.

Keep out of reach of children.

Federal law restricts this drug to use by or on the order of a licensed veterinarian.

Contains: Each tablet contains a minimum of:

Lipase	9,000 USP units
Protease	57,000 USP units
Amylase	64,000 USP units

Description: PancrePlus is a pancreatic enzyme concentrate of porcine origin. It is indicated for use as a digestive aid in replacement therapy where digestion of protein, carbohydrate and fat is inadequate due to exocrine pancreatic insufficiency.

Precautions: Discontinue use in animals with symptoms of sensitivity or allergy.

Dosage & Administration: Dose is administered before each meal. Estimate dose according to the severity of the condition and weight of the animal.

Average dose/meal:

Dogs: 2-3 tablets

Cats: 1/2-1 tablet

Storage: Store tightly closed container in a dry place at controlled room temperature.

♦ VET ONE[®] PancrePlus[™]

Powder

Pancreatic Enzyme Concentrate

For use in animals only.

Keep out of reach of children.

Federal law restricts this drug to use by or on the order of a licensed veterinarian.

Contains: Each 2.8 grams (1 teaspoon) contains a minimum of:

Lipase	71,400 USP units
Protease	388,000 USP units
Amylase	460,000 USP units

Description: PancrePlus is a pancreatic enzyme concentrate of porcine origin. It is indicated for use as a digestive aid in replacement therapy where digestion of protein, carbohydrate and fat is inadequate due to exocrine pancreatic insufficiency.

Precautions: Discontinue use in animals with symptoms of sensitivity. May cause oral mucosal irritation.

Dosage & Administration: Dose is administered before each meal. Estimate dose according to the severity of the condition and weight of the animal. PancrePlus is added to moistened food (canned or dry). Thorough mixing is necessary to bring the enzymes into close contact with the food particles. After mixing, let stand at room temperature for 15-20 minutes.

Average dose/meal:

Dogs: 3/4 to 1 teaspoon (2.8 g/teaspoon)

Cats: ¼ to ¾ teaspoon Storage: Store tightly closed container in a dry place at temperatures not exceeding 35°C (95°F).



